



**Cottage Grove Infusion**  
1515 Village Drive  
Cottage Grove, OR 97424  
Phone 541-767-5447 Fax 541-767-5399 E-Fax 541-434-3164

### **Anticoagulation Bridging Order Set (v. 02/19/2025)**

**Diagnosis/Indication for Anticoagulation (ICD-10):** \_\_\_\_\_

Follow up anti-coagulation by \_\_\_\_\_ (provider required) & \_\_\_\_\_ Anticoagulation Clinic (if applicable) after discharge from CG Infusion

.. **Bridging to warfarin (new start)**

- Start warfarin plus enoxaparin bridging tomorrow (or day after hospital discharge) per pharmacy dosing

.. **Bridging off/ on for procedure**

**Date of Procedure:** \_\_\_\_\_ **Type of procedure:** \_\_\_\_\_

Take last dose of warfarin 5 days before procedure, or on \_\_\_\_\_

Begin enoxaparin per pharmacy dosing 4 days before procedure, or on \_\_\_\_\_

.. On day after procedure, resume warfarin plus enoxaparin bridging per pharmacy dosing (or day after hospital discharge)

.. Patient will NOT return to CG Infusion for bridging after procedure

■ **Discontinuation anticoagulation bridging order set when criteria met**

- Mechanical MVR, Mechanical AVR plus a-fib: overlap for at least 4 days & INR 2.5- 3.5 for at least 1 day
- PE, DVT (new or history): overlap for at least 5 days & INR 2-3 for at least 2 days
- All other indications: overlap for at least 4 days & INR 2-3 for at least 1 day

**Labs:**

- INR, CBC no differential, BMP on day 1 (if not done in last 7 days), then per pharmacy protocol

**Nursing communications:**

- Vital signs: as needed

**Emergency Medications:**

- Diphenhydramine (BENADRYL) 25 to 50 mg IV as needed for mild to moderate drug reactions (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritic). Administer 25 mg IV once, if reaction does not resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg and contact provider.
- MethylPREDNISolone sodium succinate (Solu-MEDROL) 125 mg IV once as needed for shortness of breath, continued symptoms of mild to moderate drug reaction (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritic) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl). Contact provider if given.
- Epinephrine 0.3 mg IM once for anaphylaxis. If reaction does not resolve in 3 minutes may repeat 0.3 mg IM dose for a total of 0.6 mg. Avoid use of hand, foot, leg veins in elderly patient and those with occlusive vascular disease. Contact provider if given.
- Famotidine (PEPCID) 20 mg IV once as needed for infusion/ allergic reaction.
- Emergency medications can be given IM if IV route unavailable

Patient name: \_\_\_\_\_

Provider printed name: \_\_\_\_\_

DOB: \_\_\_\_\_

Provider signature: \_\_\_\_\_

Height \_\_\_\_\_ Weight \_\_\_\_\_

Date: \_\_\_\_\_ Time: \_\_\_\_\_