



Riverbend Medical Center Regional Infusion Center (RIC)
3377 Riverbend Drive Suite 502/510
Springfield, Oregon 97477
Phone 541-222-6280 Fax 541-349-8006

Epoetin Alfa or Biosimilar (EPOGEN, PROCRIT, RETACRIT) INFUSION (v. 11/14/2024)

Diagnosis/Indication (ICD-10): _____


Medications:

- .. Epoetin 20,000 units subcutaneous every 2 weeks. On follow up labs, if hemoglobin less than 10 g/dL but increasing continue current dose, if hemoglobin less than 10 g/dL and stable or decreasing increase dose by 25%. On follow up labs, if hemoglobin 10 -10.9 g/dL, continue current dose. On follow up labs, if hemoglobin 11-11.9 g/dL, decrease dose by 10%. On follow up labs, if hemoglobin greater than 12 g/dL, hold Epoetin dose and restart at 50% dose when hemoglobin less than 10 g/dL. (max dose 60,000 units)
- .. Epoetin fixed dose _____ units subcutaneous every _____ weeks (max dose 60,000 units)

Labs:

-  Hemoglobin every _____ week
-  Iron deficiency panel (including ferritin) every 12 weeks

Nursing communications:

-  Vital signs: Initial and as needed
-  Notify provider if blood pressure is greater than 170/100
-  Prior to first dose ensure hemoglobin is less than 10 gm/dL
-  Contact provider if dose is 60,000 units and no response
-  Hold and contact provider if hemoglobin is greater than _____ (default is 11 gm/dL)
-  If labs not resulted at time of appointment, run hemoglobin on HemoCue

Emergency Medications:

-  Diphenhydramine (BENADRYL) 25 to 50 mg IV as needed for mild to moderate drug reactions (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritic).
 -  Administer 25 mg IV once, if reaction does not resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg and contact provider.
 -  Methylprednisolone sodium succinate (Solu-MEDROL) 125 mg IV once as needed for shortness of breath, continued symptoms of mild to moderate drug reaction (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritic) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl). Contact provider if given.
-  Epinephrine 0.3 mg IM once for anaphylaxis. If reaction does not resolve in 3 minutes may repeat 0.3 mg IM dose for a total of 0.6 mg. Avoid use of hand, foot, leg veins in elderly patient and those with occlusive vascular disease. Contact provider if given.
-  Famotidine (PEPCID) 20 mg IV once as needed for infusion/ allergic reaction.
-  Emergency medications can be given IM if IV route unavailable

Patient name _____

Provider printed name: _____

DOB _____

Provider signature: _____

Height _____ Weight _____

Date: _____ Time: _____