

PENICILLIN INFUSION

Columbia Network Infusion Centers

 Select Location: Southwest/Vancouver St John/Longview Start Date: _____

Diagnosis/Indication: _____

ICD-10 Code(s): _____ Authorization Number: _____

Patient Vitals: Height _____ Weight _____

Provider Communication Orders

To reduce delays in treatment and phone calls to your office, you may participate in the PeaceHealth formulary process by selecting this option. A clinic pharmacist will adjust orders according to PeaceHealth site approved policies and procedures.

I agree to utilize the PeaceHealth policies & procedures that have been reviewed by the Pharmacy & Therapeutics Committee and authorized by the Medical Executive Committee of PeaceHealth. This agreement will be issued for the duration of active orders contained within this treatment plan.

Labs	Interval
<input type="radio"/> Complete Blood Count W/ Automated Differential	Frequency _____
<input type="radio"/> Comprehensive Metabolic Panel	Frequency _____
<input type="radio"/> C-Reactive Protein	Frequency _____
<input type="radio"/> Sedimentation Rate, Westergren	Frequency _____

Supportive Care	Interval
<input type="radio"/> PENICILLIN G CADD INFUSION (2 MU Q4H)	Route: IV
<input type="radio"/> PENICILLIN G CADD INFUSION (3 MU Q4H)	Route: IV
<input type="radio"/> PENICILLIN G CADD INFUSION (4 MU Q4H)	Route: IV
<input type="radio"/> PENICILLIN G 20 MU DAILY CONTINUOUS CADD INFUSION	Route: IV

DURATION / END DATE: _____

Nursing Orders	Interval
<input checked="" type="checkbox"/> Nursing Communication <i>At the end of treatment, Contact MD/provider to address removal of PICC line.</i>	Every Visit
<input checked="" type="checkbox"/> Nursing Communication <i>Discontinue therapy plan after treatment complete.</i>	Every Visit

Vascular Access (single select)	Interval
<input type="radio"/> IV - Insert peripheral IV - sodium chloride (NS) flush 10 mL 10 mL As Needed for Line Care	Every Visit

Provider Signature

Date

Time

Provider's Printed Name:

Place Patient Label Here

Page 1 of 2

EHR5048-DT (12/08/2024)

PENICILLIN INFUSION

- Central line (non-PICC) Every Visit
 - Access vascular device and confirm patency
 - Initiate Central line (non-PICC) maintenance protocol
 - sodium chloride (NS) flush 10 mL
 - 10 mL As Needed for Line Care prior to medication administration PRN
 - sodium chloride (NS) flush 10 mL
 - 10 mL As Needed for Line Care post medication administration PRN
 - sodium chloride (NS) flush 20 mL
 - 20 mL As Needed for Line Care post lab draw PRN
 - sodium chloride (NS) flush 10 mL
 - 10 mL for Line Care at discharge and de-access every visit
 - heparin 100 units/mL 5 mL
 - 5 mL for de-access PRN
 - alteplase (CATHFLO) injection 2 mg
 - Other, Starting when released, For 2 dose, PRN, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded
- PICC line Every Visit
 - Access vascular device and confirm patency
 - Initiate PICC maintenance protocol
 - Nursing Communication
 - Change PICC line dressing weekly and PRN
 - sodium chloride (NS) flush 10 mL
 - 10 mL As Needed for Line Care prior to medication administration PRN
 - sodium chloride (NS) flush 10 mL
 - 10 mL As Needed for Line Care post medication administration PRN
 - sodium chloride (NS) flush 20 mL
 - 20 mL As Needed for Line Care post lab draw PRN
 - alteplase (CATHFLO) injection 2 mg
 - Other, Starting when released, For 2 dose, PRN, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

PRN Medications	Interval
○ sodium chloride (NS) flush 10 mL 10 mL As Needed for Line Care	PRN Route: IV
○ sodium chloride 0.9 % bolus 500 mL 500 mL Once for Line Care, Starting when released, For 1 dose, Run at TKO (25ml/hr)	PRN Route: IV

Provider Signature

Date

Time

Provider's Printed Name:

Place Patient Label Here

Page 2 of 2

EHR5048-DT (12/08/2024)

PENICILLIN INFUSION