



Riverbend Medical Center Regional Infusion Center (RIC)
3377 Riverbend Drive Suite 502/510
Springfield, Oregon 97477
Phone 541-222-6280 Fax 541-349-8006

Risankizumab-RZAA (SKYRIZI) INFUSION Order Set (v. 10/4/2024)

Diagnosis/Indication (ICD-10): _____

* Prescribing information recommends screening for latent infections. Provider should monitor labs for hepatotoxicity

Medications:

- .. Risankizumab-RZAA 600 mg IVPB infused over 1 hour at weeks 0, 4 and 8 (Crohn's disease)
- .. Risankizumab-RZAA 1200 mg IVPB infused over 2 hours at weeks 0, 4 and 8 (Ulcerative colitis)

Nursing communications:

- Vital signs: Initial, every 30 minutes during infusion and as needed
- Hold dose and contact provider for signs of active infection
- Following induction 3 doses future therapy will be done at home

Access:

- .. Insert peripheral IV
 - Every visit, remove after IV administration complete
- .. Access & Use Central Line/ CVAD
 - Initiate Central Line (Non-PICC) Maintenance Protocol
 - Heparin, porcine (PF) 100 unit/mL flush 5 mL as needed for Port-a-Cath line care
 - Alteplase (Cathflo) 2 mg as needed for occluded catheter. For clearing central line catheter- retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded
- .. Access & Use PICC
 - Initiate PICC Maintenance Protocol
 - Normal saline flush 3 mL as needed for PICC/ Hickman line care
 - Alteplase (Cathflo) 2 mg as needed for occluded catheter. For clearing central line catheter- retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

Emergency Medications:

- Diphenhydramine (BENADRYL) 25 to 50 mg IV as needed for mild to moderate drug reactions (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>/= 20 points in SBP), nausea, urticaria, chills, pruritic). Administer 25 mg IV once, if reaction does not resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg and contact provider.
- MethylPREDNISolone sodium succinate (Solu-MEDROL) 125 mg IV once as needed for shortness of breath, continued symptoms of mild to moderate drug reaction (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>/= 20 points in SBP), nausea, urticaria, chills, pruritic) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl). Contact provider if given.

Epinephrine 0.3 mg IM once for anaphylaxis. If reaction does not resolve in 3 minutes may repeat 0.3 mg IM dose for a total of 0.6 mg. Avoid use of hand, foot, leg veins in elderly patient and those with occlusive vascular disease. Contact provider if given.

- Famotidine (PEPCID) 20 mg IV once as needed for infusion/ allergic reaction.

Patient name: _____

Provider printed name: _____

DOB: _____

Provider signature: _____

Height _____ Weight _____

Date: _____ Time: _____