



**Riverbend Medical Center Regional Infusion Center (RIC)**  
**3377 Riverbend Drive Suite 502/510**  
**Springfield, Oregon 97477**  
**Phone 541-222-6280 Fax 541-349-8006**

**USTEKINUMAB (Stelara) (v. 05/24/2024)**

**Diagnosis/Indication (ICD-10):** \_\_\_\_\_

\* Please send copy of recent (within last 30 days) CBC with differential, CMP, CRP, ESR and Quantiferon gold prior with this order

**Medication:**

- .. Usekinumab 260 mg IV x 1 (patient weight < 55kg) loading dose
- .. Usekinumab 390 mg IV x 1 (patient weight 55 - 85kg) loading dose
- .. Usekinumab 520 mg IV x 1 (patient weight > 85kg) loading dose
- .. Acetaminophen (TYLENOL) 650 mg PO once 30 minutes prior to infusion
- .. Diphenhydramine (Benadryl) 25 mg PO once 30 minutes prior to infusion

**Nursing communications:**

- n Vital signs: Initial, and as needed. Call provider for systolic BP <80 or >200 mmHg, pulse <50 or >130 bpm, temp > 38.3 C.
- .. Observe patient 30 minutes after infusion, if stable then discharge patient home

**Access:**

- .. Insert peripheral IV
  - Every visit, remove after IV administration complete
- .. Access & Use Central Line/ CVAD
  - Initiate Central Line (Non-PICC) Maintenance Protocol
  - Heparin, porcine (PF) 100 unit/mL flush 5 mL as needed for Port-a-Cath line care
  - Alteplase (Cathflo) 2 mg as needed for occluded catheter. For clearing central line catheter- retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded
- .. Access & Use PICC
  - Initiate PICC Maintenance Protocol
  - Normal saline flush 3 mL as needed for PICC/ Hickman line care
  - Alteplase (Cathflo) 2 mg as needed for occluded catheter. For clearing central line catheter- retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

**Emergency Medications:**

- n Diphenhydramine (BENADRYL) 25 to 50 mg IV as needed for mild to moderate drug reactions (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>/= 20 points in SBP), nausea, urticaria, chills, pruritic).
  - Administer 25 mg IV once, if reaction does not resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg and contact provider.
- n MethylPREDNISolone sodium succinate (Solu-MEDROL) 125 mg IV once as needed for shortness of breath, continued symptoms of mild to moderate drug reaction (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>/= 20 points in SBP), nausea, urticaria, chills, pruritic) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl). Contact provider if given.
- n Epinephrine 0.3 mg IM once for anaphylaxis. If reaction does not resolve in 3 minutes may repeat 0.3 mg IM dose for a total of 0.6 mg. Avoid use of hand, foot, leg veins in elderly patient and those with occlusive vascular disease. Contact provider if given.
- n Famotidine (PEPCID) 20 mg IV once as needed for infusion/ allergic reaction.
- n May give emergency medications IM if IV route unavailable

Patient name: \_\_\_\_\_

Provider printed name: \_\_\_\_\_

DOB: \_\_\_\_\_

Provider signature: \_\_\_\_\_

Height \_\_\_\_\_ Weight \_\_\_\_\_

Date: \_\_\_\_\_ Time: \_\_\_\_\_